

Dechant, Pierre-Philippe ORCID:

<https://orcid.org/0000-0002-4694-4010> (2017) A Clifford algebraic approach to reflection groups and root systems. In: Yau Institute Seminar in Geometry and Physics, 10th August 2017, Yau Mathematical Sciences Center (YMSC), Tsinghua, China. (Unpublished)

Downloaded from: <http://ray.yorks.ac.uk/id/eprint/4028/>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repository Policy Statement](#)

# RaY

Research at the University of York St John

For more information please contact RaY at [ray@yorks.ac.uk](mailto:ray@yorks.ac.uk)



UNIVERSITY  
*of York*



清華大學

Tsinghua University


# A Clifford algebraic approach to reflection groups and root systems

Pierre-Philippe Dechant

Mathematics Department, University of York

Yau Institute Seminar in Geometry and Physics  
August 10th, 2017

# Reflection groups: a new approach





$$s_{\alpha}(v) = v - 2 \frac{(v|\alpha)}{(\alpha|\alpha)} \alpha =$$

$$-\alpha v \alpha = -(-\alpha)v(-\alpha)$$

- Work at the level of **root systems** (which define reflection groups)
- Interested in **non-crystallographic** root systems e.g. viruses, fullerenes etc. But: no Lie algebra, so conventionally less studied
- **Clifford algebra** is a uniquely suitable framework for reflection groups/root systems: **reflection formula**, spinor **double covers**, **complex/quaternionic quantities** arising as **geometric objects**

# Main results

- Framework for reflection, conformal, modular and braid groups
- New view on the geometry of the Coxeter plane
- Induction of exceptional root systems and ADE from Platonic symmetries
- Naturally defines a range of representations



# Platonic Solids



Platonic Solid	Group	root system
Tetrahedron	$A_3$ $A_1^3$	Cuboctahedron Octahedron
Octahedron Cube	$B_3$	Cuboctahedron + Octahedron
Icosahedron Dodecahedron	$H_3$	Icosidodecahedron

- Platonic Solids have been known for millennia

# Platonic Solids



$A_1^3$

$A_3$

$B_3$





$H_3$

Platonic Solid	Group	root system
Tetrahedron	$A_3$ $A_1^3$	Cuboctahedron Octahedron
Octahedron Cube	$B_3$	Cuboctahedron + Octahedron
Icosahedron Dodecahedron	$H_3$	Icosidodecahedron

- Platonic Solids have been known for millennia
- Described by Coxeter groups

## 4D analogues of the Platonic Solids

- The 16-cell, 24-cell, 24-cell and dual 24-cell, the 600-cell and the 120-cell
- In higher dimensions there are **only** hypersimplices and hypercubes/octahedra ( $A_n$  and  $B_n$ )



# Platonic Solids



$A_1^3$

$A_1^4$

$A_3$

$D_4$

$B_3$

$F_4$

$H_3$

$H_4$


- **Abundance** of 4D root systems – **exceptional**
- Concatenating 3D reflections gives 4D **Clifford** spinors (**binary polyhedral groups**)
- These **induce 4D root systems**  

$$R = a_0 + a_1 e_2 e_3 + a_2 e_3 e_1 + a_3 e_1 e_2 \Rightarrow$$

$$R\tilde{R} = a_0^2 + a_1^2 + a_2^2 + a_3^2$$
- This construction accidental to 3D perhaps explains the unusual abundance of 4D root systems



# Root systems



Root system  $\Phi$ : set of vectors  $\alpha$  in a vector space with an inner product such that




1.  $\Phi \cap \mathbb{R}\alpha = \{-\alpha, \alpha\} \quad \forall \alpha \in \Phi$
2.  $s_\alpha \Phi = \Phi \quad \forall \alpha \in \Phi$

reflection groups

$$s_\alpha : v \rightarrow s_\alpha(v) = v - 2 \frac{(v|\alpha)}{(\alpha|\alpha)} \alpha$$



Simple roots: express every element of  $\Phi$  via a  $\mathbb{Z}$ -linear combination with coefficients of the same sign.

# The Icosahedron




- Rotational icosahedral group is  $I = A_5$  of order 60
- Full icosahedral group is  $H_3$  of order 120 (including reflections/inversion); generated by the root system icosidodecahedron

# Non-crystallographic Coxeter groups $H_2 \subset H_3 \subset H_4$



$$A = \begin{pmatrix} 2 & -\tau \\ -\tau & 2 \end{pmatrix}$$



$$A = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -\tau \\ 0 & -\tau & 2 \end{pmatrix}$$



$$A = \begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -\tau \\ 0 & 0 & -\tau & 2 \end{pmatrix}$$

$H_2 \subset H_3 \subset H_4$ : 10, 120, 14,400 elements, the only Coxeter groups that generate **rotational symmetries of order 5**.

Linear combinations now in the **extended integer ring**

$$\mathbb{Z}[\tau] = \{a + \tau b \mid a, b \in \mathbb{Z}\}$$

**golden ratio**

$$\tau = \frac{1}{2}(1 + \sqrt{5}) = 2 \cos \frac{\pi}{5}$$


$$x^2 = x + 1$$

$$\tau' = \sigma = \frac{1}{2}(1 - \sqrt{5}) = 2 \cos \frac{2\pi}{5}$$

$$\tau + \sigma = 1, \tau\sigma = -1$$

# Cartan-Dynkin diagrams

Coxeter-Dynkin diagrams: **node** = **simple** root, no link = roots orthogonal i.e. **angle**  $\frac{\pi}{2}$ , simple link = roots at **angle**  $\frac{\pi}{3}$ , link with label  $m$  = **angle**  $\frac{\pi}{m}$ .



- 1 Polyhedral groups, Platonic solids and root systems
- 2 Reflection groups with Clifford algebras
  - A Clifford way of doing orthogonal transformations
  - The geometry of the Coxeter plane
  - Root system induction and ADE correspondences
  - Representations from multivector groups
  - Conformal, modular and braid groups
- 3 Conclusions

# Clifford Algebra and orthogonal transformations

- **Geometric Product** for two vectors  $ab \equiv a \cdot b + a \wedge b$
- **Inner product** is symmetric part  $a \cdot b = \frac{1}{2}(ab + ba)$
- Reflecting  $a$  in  $n$  is given by  $a' = a - 2(a \cdot n)n = -nan$  ( $n$  and  $-n$  **doubly cover** the same reflection)
- Via **Cartan-Dieudonné** theorem any orthogonal transformation can be written as **successive reflections**, which are **doubly covered** by Clifford versors/pinors  $A$

$$x' = \pm n_1 n_2 \dots n_k x n_k \dots n_2 n_1 =: \pm A x \tilde{A}$$

## Clifford Algebra of 3D: the relation with 4D and 8D

- Clifford (Pauli) algebra in 3D is

$$\underbrace{\{1\}}_{1 \text{ scalar}} \quad \underbrace{\{e_1, e_2, e_3\}}_{3 \text{ vectors}} \quad \underbrace{\{e_1 e_2, e_2 e_3, e_3 e_1\}}_{3 \text{ bivectors}} \quad \underbrace{\{I \equiv e_1 e_2 e_3\}}_{1 \text{ trivector}}$$


- We can multiply together root vectors in this algebra  $\alpha_i \alpha_j \dots$
- A general element has 8 components: 8D
- even products (rotations/spinors) have four components:

$$R = a_0 + a_1 e_2 e_3 + a_2 e_3 e_1 + a_3 e_1 e_2 \Rightarrow R \tilde{R} = a_0^2 + a_1^2 + a_2^2 + a_3^2$$

- So behaves as a 4D Euclidean object – inner product

$$(R_1, R_2) = \frac{1}{2}(R_2 \tilde{R}_1 + R_1 \tilde{R}_2)$$

## Spinors from reflections: easy example



- The 6 **roots**  $(\pm 1, 0, 0)$  and permutations in  $A_1 \times A_1 \times A_1$
- $\{\pm e_1, \pm e_2, \pm e_3\}$  generate **group of 8 spinors**  
 $\{\pm 1, \pm e_1 e_2, \pm e_2 e_3, \pm e_3 e_1\}$
- This is a **discrete spinor group** isomorphic to the **quaternion** group  $Q$ .



## Pinors from reflections: easy example

$$\underbrace{\{\pm 1\}}_{1 \text{ scalar}} \quad \underbrace{\{\pm e_1, \pm e_2, \pm e_3\}}_{3 \text{ vectors}} \quad \underbrace{\{\pm e_1 e_2, \pm e_2 e_3, \pm e_3 e_1\}}_{3 \text{ bivectors}} \quad \underbrace{\{\pm I \equiv e_1 e_2 e_3\}}_{1 \text{ trivector}}$$

- The **pin group** also of course contains  $\boxed{\pm e_1, \pm e_2, \pm e_3}$  and  $\boxed{\pm e_1 e_2 e_3}$
- So total pin group is a group of **order 16**
- Since  $\boxed{e_1, e_2, e_3}$  generate the **inversion**  $e_1 e_2 e_3$ , actually the 8 elements in the even subalgebra and the other 8 elements in the other 4D can be '**Hodge**' dualised
- So **when the group contains the inversion**  $\text{Pin} = \text{Spin} \times \mathbb{Z}_2$

## Spinors from reflections: icosahedral case

- The  $H_3$  root system has 30 **roots** e.g. simple roots

$$\alpha_1 = e_2, \alpha_2 = -\frac{1}{2}((\tau-1)e_1 + e_2 + \tau e_3) \text{ and } \alpha_3 = e_3.$$

- Subgroup of **rotations**  $A_5$  of order **60** is doubly covered by **120**

spinors of the form  $\alpha_1 \alpha_2 = -\frac{1}{2}(1 - (\tau-1)e_1 e_2 + \tau e_2 e_3),$

$$\alpha_1 \alpha_3 = e_2 e_3 \text{ and } \alpha_2 \alpha_3 = -\frac{1}{2}(\tau - (\tau-1)e_3 e_1 + e_2 e_3).$$

- The inclusion of the  $H_3$  **inversion doubles** this

## Polyhedral groups as multivector groups

Group	Discrete subgroup	Order	Action Mechanism
$SO(3)$	rotational (chiral)	$ G $	$x \rightarrow \tilde{R}xR$
$O(3)$	reflection (full/Coxeter)	$2 G $	$x \rightarrow \pm \tilde{A}xA$
$Spin(3)$	binary	$2 G $	$(R_1, R_2) \rightarrow R_1 R_2$
$Pin(3)$	pinory (?)	$4 G $	$(A_1, A_2) \rightarrow A_1 A_2$

- e.g. the **chiral icosahedral** group has 60 elements, encoded by 120 spinors, which form the **binary icosahedral** group
- together with the **inversion/pseudoscalar**  $I$  this gives 60 rotations and 60 rotoinversions, i.e. the **full icosahedral** group  $H_3$  in 120 elements doubly covered by 240 pinors

## Some Group Theory: chiral, full, binary, pin

- Easy to calculate conjugacy classes etc
- Chiral (binary) polyhedral groups have irreps
- tetrahedral (12/24):  $1, 1', 1'', 2_s, 2'_s, 2''_s, 3$
- octahedral (24/48):  $1, 1', 2, 2_s, 2'_s, 3, 3', 4_s$
- icosahedral (60/120):  $1, 2_s, 2'_s, 3, \bar{3}, 4, 4_s, 5, 6_s$
- All binary are discrete subgroups of  $SU(2)$  and all thus have a  $2_s$  spinor irrep
- Connection with Trinities and the McKay correspondence

# Tetrahedral group $A_3$ : rotational group $\tilde{R} \times R$

Simple roots for  $A_3$ :

$$\alpha_1 = \frac{1}{\sqrt{2}}(e_2 - e_1), \quad \alpha_2 = \frac{1}{\sqrt{2}}(e_3 - e_2) \text{ and } \alpha_3 = \frac{1}{\sqrt{2}}(e_1 + e_2)$$

Conj. Class	Distinct rotations given by two spinors each ( $\pm$ )	
1	$\pm 1$	
4	$\pm \frac{1}{2}(1 - e_1 e_2 + e_2 e_3 - e_3 e_1),$	$\pm \frac{1}{2}(1 - e_1 e_2 - e_2 e_3 + e_3 e_1),$
	$\pm \frac{1}{2}(1 + e_1 e_2 - e_2 e_3 - e_3 e_1),$	$\pm \frac{1}{2}(1 + e_1 e_2 + e_2 e_3 + e_3 e_1)$
$4^{-1}$	$\pm \frac{1}{2}(1 + e_1 e_2 - e_2 e_3 + e_3 e_1),$	$\pm \frac{1}{2}(1 + e_1 e_2 + e_2 e_3 - e_3 e_1),$
	$\pm \frac{1}{2}(1 - e_1 e_2 + e_2 e_3 + e_3 e_1),$	$\pm \frac{1}{2}(1 - e_1 e_2 - e_2 e_3 - e_3 e_1)$
3	$\pm e_1 e_2, \quad \pm e_2 e_3, \quad \pm e_3 e_1$	

# Tetrahedral group $A_3$ : spinor group $R_1 R_2$

Conjugacy Class	Group elements
$1$	$1$
$1_-$	$-1$
$4$	$\frac{1}{2}(1 - e_1 e_2 + e_2 e_3 - e_3 e_1), \quad \frac{1}{2}(1 - e_1 e_2 - e_2 e_3 + e_3 e_1),$ $\frac{1}{2}(1 + e_1 e_2 - e_2 e_3 - e_3 e_1), \quad \frac{1}{2}(1 + e_1 e_2 + e_2 e_3 + e_3 e_1)$
$4_-$	$-\frac{1}{2}(1 - e_1 e_2 + e_2 e_3 - e_3 e_1), \quad -\frac{1}{2}(1 - e_1 e_2 - e_2 e_3 + e_3 e_1),$ $-\frac{1}{2}(1 + e_1 e_2 - e_2 e_3 - e_3 e_1), \quad -\frac{1}{2}(1 + e_1 e_2 + e_2 e_3 + e_3 e_1)$
$4^{-1}$	$\frac{1}{2}(1 + e_1 e_2 - e_2 e_3 + e_3 e_1), \quad \frac{1}{2}(1 + e_1 e_2 + e_2 e_3 - e_3 e_1),$ $\frac{1}{2}(1 - e_1 e_2 + e_2 e_3 + e_3 e_1), \quad \frac{1}{2}(1 - e_1 e_2 - e_2 e_3 - e_3 e_1)$
$4_-^{-1}$	$-\frac{1}{2}(1 + e_1 e_2 - e_2 e_3 + e_3 e_1), \quad -\frac{1}{2}(1 + e_1 e_2 + e_2 e_3 - e_3 e_1),$ $-\frac{1}{2}(1 - e_1 e_2 + e_2 e_3 + e_3 e_1), \quad -\frac{1}{2}(1 - e_1 e_2 - e_2 e_3 - e_3 e_1)$
$6$	$\pm e_1 e_2, \quad \pm e_2 e_3, \quad \pm e_3 e_1$

# Tetrahedral group $A_3$ : pin group $A_1A_2$

Conjugacy Class	Group elements
1	1
1 <sub>-</sub>	-1
8 <sub>+</sub>	$\frac{1}{2}(1 - e_1e_2 + e_2e_3 - e_3e_1), \quad \frac{1}{2}(1 - e_1e_2 - e_2e_3 + e_3e_1),$ $\frac{1}{2}(1 + e_1e_2 - e_2e_3 - e_3e_1), \quad \frac{1}{2}(1 + e_1e_2 + e_2e_3 + e_3e_1),$ $\frac{1}{2}(1 + e_1e_2 - e_2e_3 + e_3e_1), \quad \frac{1}{2}(1 + e_1e_2 + e_2e_3 - e_3e_1),$ $\frac{1}{2}(1 - e_1e_2 + e_2e_3 + e_3e_1), \quad \frac{1}{2}(1 - e_1e_2 - e_2e_3 - e_3e_1)$
8 <sub>-</sub>	$-\frac{1}{2}(1 - e_1e_2 + e_2e_3 - e_3e_1), \quad -\frac{1}{2}(1 - e_1e_2 - e_2e_3 + e_3e_1),$ $-\frac{1}{2}(1 + e_1e_2 - e_2e_3 - e_3e_1), \quad -\frac{1}{2}(1 + e_1e_2 + e_2e_3 + e_3e_1),$ $-\frac{1}{2}(1 + e_1e_2 - e_2e_3 + e_3e_1), \quad -\frac{1}{2}(1 + e_1e_2 + e_2e_3 - e_3e_1),$ $-\frac{1}{2}(1 - e_1e_2 + e_2e_3 + e_3e_1), \quad -\frac{1}{2}(1 - e_1e_2 - e_2e_3 - e_3e_1)$
6	$\pm e_1e_2, \quad \pm e_2e_3, \quad \pm e_3e_1$
12	$\frac{1}{\sqrt{2}}(\pm e_1 \pm e_2), \quad \frac{1}{\sqrt{2}}(\pm e_2 \pm e_3), \quad \frac{1}{\sqrt{2}}(\pm e_3 \pm e_1)$
6 <sub>+</sub>	$\frac{1}{\sqrt{2}}(I \pm e_1), \quad \frac{1}{\sqrt{2}}(I \pm e_2), \quad \frac{1}{\sqrt{2}}(I \pm e_3)$
6 <sub>-</sub>	$-\frac{1}{\sqrt{2}}(I \pm e_1), \quad -\frac{1}{\sqrt{2}}(I \pm e_2), \quad -\frac{1}{\sqrt{2}}(I \pm e_3)$




Doubly covers  $A_3$ .

- 1 Polyhedral groups, Platonic solids and root systems
- 2 Reflection groups with Clifford algebras
  - A Clifford way of doing orthogonal transformations
  - The geometry of the Coxeter plane
  - Root system induction and ADE correspondences
  - Representations from multivector groups
  - Conformal, modular and braid groups
- 3 Conclusions



# The Coxeter Plane

- **Every** (for our purposes) Coxeter group has a Coxeter plane.
- A way to visualise Coxeter groups in any dimension by **projecting** their root system onto the Coxeter plane



# Coxeter Elements, Degrees and Exponents

- Like the symmetric group, Coxeter groups can have **invariant polynomials**. Their **degrees**  $d$  are important invariants/group characteristics.
- Turns out that actually **degrees**  $d$  are intimately related to so-called **exponents**  $m$   $m = d - 1$ .

## Coxeter Elements, Degrees and Exponents



- A **Coxeter Element** is any combination of all the simple reflections  $w = s_1 \dots s_n$ , i.e. in Clifford algebra it is encoded by the versor  $W = \alpha_1 \dots \alpha_n$  acting as  $v \rightarrow wv = \pm \tilde{W}vW$ . All such elements are conjugate and thus their **order** is invariant and called the **Coxeter number**  $h$ .
- The Coxeter element has **complex eigenvalues** of the form  $\exp(2\pi mi/h)$  where  $m$  are called **exponents**:  
$$wx = \exp(2\pi mi/h)x$$
- Standard theory **complexifies** the real Coxeter group situation in order to find **complex eigenvalues**, then takes **real sections** again.

## Coxeter Elements, Degrees and Exponents

- The Coxeter element has **complex eigenvalues** of the form  $\exp(2\pi mi/h)$  where  $m$  are called **exponents**
- Standard theory **complexifies** the real Coxeter group situation in order to find **complex eigenvalues**, then takes **real** sections again.
- In particular, **1** and  **$h-1$**  are always exponents
- Turns out that actually **exponents and degrees** are intimately related ( **$m = d-1$** ). The construction is slightly roundabout but uniform, and uses the **Coxeter plane**.

# The Coxeter Plane

- In particular, can show **every** (for our purposes) Coxeter group has a Coxeter plane
- Existence relies on the fact that all groups in question have **tree-like Dynkin diagrams**, and thus admit an **alternate colouring**
- Essentially just gives **two sets of mutually commuting generators**



# The Coxeter Plane

- Existence relies on the fact that all groups in question have **tree-like Dynkin diagrams**, and thus admit an alternate colouring
- Essentially just gives **two sets of orthogonal = mutually commuting generators but anticommuting root vectors**  $\alpha_w$  and  $\alpha_b$  (duals  $\omega$ )
- Cartan matrices are positive definite, and thus have a **Perron-Frobenius** (all positive) eigenvector  $\lambda_i$ .
- Take **linear combinations** of components of this eigenvector as coefficients of two vectors from the orthogonal sets  
$$v_w = \sum \lambda_w \omega_w \text{ and } v_b = \sum \lambda_b \omega_b$$
- Their **outer product/Coxeter plane bivector**  $B_C = v_b \wedge v_w$  describes an **invariant plane** where  $w$  acts by rotation by  $2\pi/h$ .

## Clifford Algebra and the Coxeter Plane – 2D case

$$I_2(n) \quad \circ \xrightarrow{n} \circ$$

- For  $I_2(n)$  take  $\alpha_1 = e_1, \alpha_2 = -\cos \frac{\pi}{n} e_1 + \sin \frac{\pi}{n} e_2$

- So **Coxeter versor** is just

$$W = \alpha_1 \alpha_2 = -\cos \frac{\pi}{n} + \sin \frac{\pi}{n} e_1 e_2 = -\exp \left( -\frac{\pi I}{n} \right)$$

- In Clifford algebra it is therefore immediately obvious that the action of the  $I_2(n)$  Coxeter element is described by a versor (here a rotor/spinor) that encodes **rotations in the  $e_1 e_2$ -Coxeter-plane** and yields  **$h = n$**  since trivially  $W^n = (-1)^{n+1}$  yielding  $w^n = 1$  via  $wv = \tilde{W}vW$ .

## Clifford Algebra and the Coxeter Plane – 2D case

- Coxeter versor  $W = -\cos \frac{\pi}{n} + \sin \frac{\pi}{n} e_1 e_2 = -\exp \left( -\frac{\pi I}{n} \right)$

- $I = e_1 e_2$  anticommutes with both  $e_1$  and  $e_2$  such that sandwiching formula becomes

$$v \rightarrow wv = \tilde{W} v W = \tilde{W}^2 v = \exp \left( \pm \frac{2\pi I}{n} \right) v \text{ immediately}$$

yielding the standard result for the complex eigenvalues in real Clifford algebra without any need for artificial complexification

- The Coxeter plane bivector  $B_C = e_1 e_2 = I$  gives the complex structure
- The Coxeter plane bivector  $B_C$  is invariant under the Coxeter versor  $\tilde{W} B_C W = \pm B_C$ .



## Clifford algebra: no need for complexification

- Turns out in Clifford algebra we can **factorise**  $W$  into **orthogonal** (commuting/anticommuting) components

$$W = \alpha_1 \dots \alpha_n = W_1 \dots W_n \text{ with } W_i = \exp(\pi m_i l_i / h)$$

- Here,  $l_i$  is a bivector describing a **plane** with  $l_i^2 = -1$

- For  $v$  **orthogonal to the plane** described by  $l_i$  we have

$$v \rightarrow \tilde{W}_i v W_i = \tilde{W}_i W_i v = v \text{ so cancels out}$$

- For  $v$  **in the plane** we have

$$v \rightarrow \tilde{W}_i v W_i = \tilde{W}_i^2 v = \exp(2\pi m_i l_i / h) v$$

- Thus if we **decompose**  $W$  into **orthogonal eigenspaces**, in the eigenvector equation all orthogonal bits cancel out and one gets the complex eigenvalue from the respective eigenspace

# Clifford algebra: no need for complexification

- For  $v$  in the plane we have

$$v \rightarrow \tilde{W}_i v W_i = \tilde{W}_i^2 v = \exp(2\pi m_i l_i / h) v$$

- So **complex eigenvalue equation** arises geometrically **without any need** for complexification
- **Different complex structures** immediately give different **eigenplanes**
- Eigenvalues/angles/**exponents** given from just factorising  $W = \alpha_1 \dots \alpha_n$
- E.g.  $H_4$  has exponents 1, 11, 19, 29 and  $W = \exp\left(\frac{\pi}{30} B_C\right) \exp\left(\frac{11\pi}{30} I B_C\right)$
- Here we have been looking for orthogonal eigenspaces, so **innocuous** – different complex structures commute
- But not in general – **naive complexification** can be misleading


## 4D case: $D_4$

- E.g.  $D_4$  has exponents 1, 3, 3, 5
- Coxeter versor decomposes into **orthogonal components**

$$W = \alpha_1 \alpha_2 \alpha_3 \alpha_4 = e_1 e_2 e_3 e_4 - e_2 e_3 - e_1 e_2 + e_1 e_3$$

$$= \frac{1}{2}(\sqrt{3} - B_C)IB_C = \exp\left(\frac{\pi}{6}B_C\right)\exp\left(\frac{3\pi}{6}IB_C\right)$$



$$B_C = 1/\sqrt{3}(e_1 + e_2 + e_3)e_4; \quad IB_C = (e_1 + e_2 - 2e_3)(e_1 - e_2)$$



## 4D case: $F_4$

- E.g.  $F_4$  has exponents 1,5,7,11
- Coxeter versor decomposes into **orthogonal components**



$$W = \alpha_1 \alpha_2 \alpha_3 \alpha_4 = \exp\left(\frac{\pi}{12} B_C\right) \exp\left(\frac{5\pi}{12} I B_C\right)$$



## 4D case: $H_4$

- E.g.  $H_4$  has exponents 1, 11, 19, 29
- Coxeter versor decomposes into orthogonal components

$$W = \alpha_1 \alpha_2 \alpha_3 \alpha_4 = \exp\left(\frac{\pi}{30} B_C\right) \exp\left(\frac{11\pi}{30} I B_C\right)$$





# Clifford Algebra and the Coxeter Plane – 4D case summary

rank 4	exponents	W-factorisation
$A_4$	1, 2, 3, 4	$W = \exp\left(\frac{\pi}{5} B_C\right) \exp\left(\frac{2\pi}{5} I B_C\right)$
$B_4$	1, 3, 5, 7	$W = \exp\left(\frac{\pi}{8} B_C\right) \exp\left(\frac{3\pi}{8} I B_C\right)$
$D_4$	1, 3, 3, 5	$W = \exp\left(\frac{\pi}{6} B_C\right) \exp\left(\frac{\pi}{2} I B_C\right)$
$F_4$	1, 5, 7, 11	$W = \exp\left(\frac{\pi}{12} B_C\right) \exp\left(\frac{5\pi}{12} I B_C\right)$
$H_4$	1, 11, 19, 29	$W = \exp\left(\frac{\pi}{30} B_C\right) \exp\left(\frac{11\pi}{30} I B_C\right)$

## 8D case: $E_8$

- E.g.  $H_4$  has exponents 1, 11, 19, 29,  $E_8$  has 1, 7, 11, 13, 17, 19, 23, 29
- Coxeter versor decomposes into **orthogonal components**



$$W = \alpha_1 \dots \alpha_8 = \exp\left(\frac{\pi}{30} B_C\right) \exp\left(\frac{7\pi}{30} B_2\right) \exp\left(\frac{11\pi}{30} B_3\right) \exp\left(\frac{13\pi}{30} B_4\right)$$



## 8D case: $E_8$

- E.g.  $H_4$  has exponents 1, 11, 19, 29,  $E_8$  has 1, 7, 11, 13, 17, 19, 23, 29
- Coxeter versor decomposes into **orthogonal components**

$$W = \alpha_1 \dots \alpha_8 = \exp\left(\frac{\pi}{30} B_C\right) \exp\left(\frac{7\pi}{30} B_2\right) \exp\left(\frac{11\pi}{30} B_3\right) \exp\left(\frac{13\pi}{30} B_4\right)$$





## Imaginary differences – different imaginaries


So what has been **gained** by this **Clifford view**?

- There are **different** entities that serve as **unit imaginaries**
- They have a **geometric** interpretation as an **eigenplane of the Coxeter element**
- These don't need to **commute** with everything like  $i$  (though they do here – at least anticommute. But that is because we looked for **orthogonal decompositions**)
- But see that in general **naive complexification** can be a dangerous thing to do – **unnecessary**, issues of **commutativity**, **confusing** different imaginaries etc



- 1 Polyhedral groups, Platonic solids and root systems
- 2 Reflection groups with Clifford algebras
  - A Clifford way of doing orthogonal transformations
  - The geometry of the Coxeter plane
  - Root system induction and ADE correspondences
  - Representations from multivector groups
  - Conformal, modular and braid groups
- 3 Conclusions

## Exceptional $E_8$ (projected into the Coxeter plane)

$E_8$  root system has 240 roots,  $H_3$  has order 120



- Order 120 group  $H_3$  doubly covered by 240 (s)pinors in 8D space
- With (somewhat counterintuitive) reduced inner product this gives the  $E_8$  root system
- $E_8$  is actually hidden within 3D geometry!



## Induction Theorem – root systems

- Induction Theorem: every 3D root system gives a 3D spinor group which gives a 4D root system.

## Induction Theorem – root systems

- Induction Theorem: every 3D root system gives a 3D spinor group which gives a 4D root system.

- Check axioms:

1.  $\Phi \cap \mathbb{R}\alpha = \{-\alpha, \alpha\} \quad \forall \alpha \in \Phi$

2.  $s_\alpha \Phi = \Phi \quad \forall \alpha \in \Phi$

## Induction Theorem – root systems


- Induction Theorem: every 3D root system gives a 3D spinor group which gives a 4D root system.
- Check axioms:
  1.  $\Phi \cap \mathbb{R}\alpha = \{-\alpha, \alpha\} \quad \forall \alpha \in \Phi$
  2.  $s_\alpha \Phi = \Phi \quad \forall \alpha \in \Phi$
- Proof: 1.  $R$  and  $-R$  are in a spinor group by construction (double cover of orthogonal transformations), 2. closure under reflections is guaranteed by the closure property of the spinor group (with a twist:  $-R_1 \tilde{R}_2 R_1$ )

## Induction Theorem – root systems

- Induction Theorem: every 3D root system gives a 3D spinor group which gives a 4D root system.
- Check axioms:
  1.  $\Phi \cap \mathbb{R}\alpha = \{-\alpha, \alpha\} \quad \forall \alpha \in \Phi$
  2.  $s_\alpha \Phi = \Phi \quad \forall \alpha \in \Phi$
- Proof: 1.  $R$  and  $-R$  are in a spinor group by construction (double cover of orthogonal transformations), 2. closure under reflections is guaranteed by the closure property of the spinor group (with a twist:  $-R_1 \tilde{R}_2 R_1$ )
- In 2D, the space of spinors is also 2D and the root systems are self-dual under an analogous construction



## Spinors from reflections: easy example



- The 6 **roots**  $(\pm 1, 0, 0)$  and permutations in  $A_1 \times A_1 \times A_1$  generate 8 **spinors**:
- $\boxed{\pm e_1, \pm e_2, \pm e_3}$  give the 8 spinors  $\boxed{\pm 1, \pm e_1 e_2, \pm e_2 e_3, \pm e_3 e_1}$
- This is a **discrete spinor group** isomorphic to the **quaternion** group  $Q$ .
- As 4D vectors these are  $(\pm 1, 0, 0, 0)$  and permutations, the 8 **roots** of  $A_1 \times A_1 \times A_1 \times A_1$  (the 16-cell).

## $H_4$ from $H_3$

- The  $H_3$  root system has 30 **roots** e.g. simple roots

$$\alpha_1 = e_2, \alpha_2 = -\frac{1}{2}((\tau-1)e_1 + e_2 + \tau e_3) \text{ and } \alpha_3 = e_3.$$

- Subgroup of **rotations**  $A_5$  of order **60** is doubly covered by **120**

spinors of the form  $\alpha_1 \alpha_2 = -\frac{1}{2}(1 - (\tau-1)e_1 e_2 + \tau e_2 e_3),$

$\alpha_1 \alpha_3 = e_2 e_3$  and  $\alpha_2 \alpha_3 = -\frac{1}{2}(\tau - (\tau-1)e_3 e_1 + e_2 e_3).$

- 





$$(\pm 1, 0, 0, 0) \text{ (8 perms)}, \quad \frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1) \text{ (16 perms)}$$

$$\frac{1}{2}(0, \pm 1, \pm \sigma, \pm \tau) \text{ (96 even perms),}$$

As **4D vectors** are the 120 roots of the  **$H_4$  root system**.

# Spinors and Polytopes

- Can reinterpret **spinors in  $\mathbb{R}^3$**  as **vectors in  $\mathbb{R}^4$**
- Give (exceptional) root systems ( $D_4, F_4, H_4$ )
- They constitute the **vertices** of the **16-cell**, **24-cell**, **24-cell** and **dual 24-cell** and the **600-cell**
- These are 4D analogues of the **Platonic Solids**. **Strange symmetries** better understood in terms of **3D spinors**



# Trinity of 4D Exceptional Root Systems

- **Exceptional** phenomena:  $D_4$  (**triality**, important in string theory),  $F_4$  (**largest lattice symmetry** in 4D),  $H_4$  (**largest non-crystallographic symmetry**); **Exceptional**  $D_4$  and  $F_4$  arise from **series**  $A_3$  and  $B_3$ ;  $A_1 \times I_2(n) \rightarrow I_2(n) \times I_2(n)$

rank-3 group	diagram	binary	rank-4 group	diagram
$A_1 \times A_1 \times A_1$		$Q$	$A_1 \times A_1 \times A_1 \times A_1$	
$A_3$		$2T$	$D_4$	
$B_3$		$2O$	$F_4$	
$H_3$		$2I$	$H_4$	

# Arnold's indirect connection between Trinities $(A_3, B_3, H_3)$ and $(D_4, F_4, H_4)$

- **Arnold** had noticed a handwavey connection:
- Decomposition of 3D groups in terms of number of **Springer cones** matches what are essentially the **exponents** of the 4D groups:
- $A_3$ :  $24 = 2(1 + 3 + 3 + 5) - D_4$ :  $(1, 3, 3, 5)$
- $B_3$ :  $48 = 2(1 + 5 + 7 + 11) - F_4$ :  $(1, 5, 7, 11)$
- $H_3$ :  $120 = 2(1 + 11 + 19 + 29) - H_4$ :  $(1, 11, 19, 29)$

## Arnold's indirect connection between Trinities

rank 4	exponents	W-factorisation
$A_4$	1, 2, 3, 4	$W = \exp\left(\frac{\pi}{5} B_C\right) \exp\left(\frac{2\pi}{5} I B_C\right)$
$B_4$	1, 3, 5, 7	$W = \exp\left(\frac{\pi}{8} B_C\right) \exp\left(\frac{3\pi}{8} I B_C\right)$
$D_4$	1, 3, 3, 5	$W = \exp\left(\frac{\pi}{6} B_C\right) \exp\left(\frac{\pi}{2} I B_C\right)$
$F_4$	1, 5, 7, 11	$W = \exp\left(\frac{\pi}{12} B_C\right) \exp\left(\frac{5\pi}{12} I B_C\right)$
$H_4$	1, 11, 19, 29	$W = \exp\left(\frac{\pi}{30} B_C\right) \exp\left(\frac{11\pi}{30} I B_C\right)$

The **remaining cases** in the root system induction construction work the same way, not just this Trinity! So more general correspondence:

$$(A_1 \times I_2(n), A_3, B_3, H_3) \rightarrow (I_2(n) \times I_2(n), D_4, F_4, H_4)$$

# The countably infinite family $I_2(n)$ and Arnold's construction

- For  $A_1^3$  can see immediately  $8 = 2(1 + 1 + 1 + 1)$
- Simple roots  $\alpha_1 = e_1, \alpha_2 = e_2, \alpha_3 = e_3, \alpha_4 = e_4$  give
$$W = e_1 e_2 e_3 e_4 = \left(\cos \frac{\pi}{2} + \sin \frac{\pi}{2} e_1 e_2\right) \left(\cos \frac{\pi}{2} + \sin \frac{\pi}{2} e_3 e_4\right) = \exp\left(\frac{\pi}{2} e_1 e_2\right) \exp\left(\frac{\pi}{2} e_3 e_4\right)$$
- Gives exponents  $(1, 1, 1, 1)$  (from  $h - 1 = 2 - 1$ )

# The countably infinite family $I_2(n)$ and Arnold's construction

- For  $A_1 \times I_2(n)$  one gets the same decomposition  
 $4n = 2(1 + (n-1) + 1 + (n-1)) = 2 \cdot 2n$
- Simple roots  $\alpha_1 = e_1$ ,  $\alpha_2 = -\cos \frac{\pi}{n} e_1 + \sin \frac{\pi}{n} e_2$ ,  $\alpha_3 = e_3$ ,  
 $\alpha_4 = -\cos \frac{\pi}{n} e_3 + \sin \frac{\pi}{n} e_4$  give  $W = \exp\left(-\frac{\pi e_1 e_2}{n}\right) \exp\left(-\frac{\pi e_3 e_4}{n}\right)$
- Gives exponents  $(1, (n-1), 1, (n-1))$



## The countably infinite family $I_2(n)$ and Arnold's construction

- So Arnold's initial hunch regarding the exponents **extends in fact to my full correspondence**
- **McKay correspondence** is a correspondence between even subgroups of  $SU(2)/\text{quaternions}$  and ADE affine Lie algebras
- In fact here get the even quaternion subgroups from 3D – **link to ADE affine Lie algebras** via McKay?

## 3D, 4D and ADE correspondences

- McKay correspondence relates even  $SU(2)$  subgroups with ADE Lie algebras ( $A_{2n-1}, D_{n+2}, E_6, E_7, E_8$ )
- Induction theorem: get these as 2D/4D root systems ( $I_2(n) \times I_2(n), D_4, F_4, H_4$ ) from 2D/3D root systems  $A_1 \times I_2(n), A_3, B_3, H_3$ )
- $(2n+2, 12, 18, 30)$  are numbers of roots, the sum of the dimensions of the irreps and the ADE Coxeter number

	4D	$G$	$\sum d_i$	ADE	$h$
				$\tilde{A}_{2n-1}$	$2n$
	$I_2(n) \times I_2(n)$	$\text{Dic}_n$	$2n+2$	$\tilde{D}_{n+2}$	$2(n+1)$
	$D_4$	$2T$	12	$\tilde{E}_6$	12
	$F_4$	$2O$	18	$\tilde{E}_7$	18
	$H_4$	$2I$	30	$\tilde{E}_8$	30

## 2D/3D, 2D/4D and ADE correspondences

- McKay correspondence relates even  $SU(2)$  subgroups with ADE Lie algebras ( $A_{2n-1}, D_{n+2}, E_6, E_7, E_8$ )
- Induction theorem: get these as 2D/4D root systems ( $I_2(n) \times I_2(n), D_4, F_4, H_4$ ) from 2D/3D root systems  $A_1 \times I_2(n), A_3, B_3, H_3$
- $(2n+2, 12, 18, 30)$  are numbers of roots, the sum of the dimensions of the irreps and the ADE Coxeter number


2D/3D	$ \Phi $	4D	$G$	$\sum d_i$	ADE	$h$
$A_1 \times I_2(n)$	$2n+2$	$I_2(n) \times I_2(n)$	$\text{Dic}_n$	$2n+2$		
$A_3$	12	$D_4$	$2T$	12		
$B_3$	18	$F_4$	$2O$	18		
$H_3$	30	$H_4$	$2I$	30		

## 2D/3D, 2D/4D and ADE correspondences

- McKay correspondence relates even  $SU(2)$  subgroups with ADE Lie algebras ( $A_{2n-1}, D_{n+2}, E_6, E_7, E_8$ )
- Induction theorem: get these as 2D/4D root systems ( $I_2(n), I_2(n) \times I_2(n), D_4, F_4, H_4$ ) from 2D/3D root systems ( $I_2(n), A_1 \times I_2(n), A_3, B_3, H_3$ )
- $(2n, 2n+2, 12, 18, 30)$  are numbers of roots, the sum of the dimensions of the irreps and the ADE Coxeter number

2D/3D	$ \Phi $	4D	$G$	$\sum d_i$	ADE	$h$
$I_2(n)$	$2n$	$I_2(n)$	$C_{2n}$	$2n$	$\tilde{A}_{2n-1}$	$2n$
$A_1 \times I_2(n)$	$2n+2$	$I_2(n) \times I_2(n)$	$\text{Dic}_n$	$2n+2$	$\tilde{D}_{n+2}$	$2(n+1)$
$A_3$	12	$D_4$	$2T$	12	$\tilde{E}_6$	12
$B_3$	18	$F_4$	$2O$	18	$\tilde{E}_7$	18
$H_3$	30	$H_4$	$2I$	30	$\tilde{E}_8$	30


# Is there a direct Platonic-ADE correspondence?



2D/3D		rot	ADE		legs
$I_2(n)$		$n$	$A_n$		$n$
$A_1 \times I_2(n)$		$2, 2, n$	$D_{n+2}$		$2, 2, n$
$A_3$		$2, 3, 3$	$E_6$		$2, 3, 3$
$B_3$		$2, 3, 4$	$E_7$		$2, 3, 4$
$H_3$		$2, 3, 5$	$E_8$		$2, 3, 5$

## A Trinity of root system ADE correspondences

- **2D/3D** root systems ( $I_2(n), A_1 \times I_2(n), A_3, B_3, H_3$ )
- **2D/4D** root systems ( $I_2(n), I_2(n) \times I_2(n), D_4, F_4, H_4$ )
- **ADE** root systems ( $A_n, D_{n+2}, E_6, E_7, E_8$ )



- 1 Polyhedral groups, Platonic solids and root systems
- 2 Reflection groups with Clifford algebras
  - A Clifford way of doing orthogonal transformations
  - The geometry of the Coxeter plane
  - Root system induction and ADE correspondences
  - Representations from multivector groups
  - Conformal, modular and braid groups
- 3 Conclusions

## Polyhedral groups as multivector groups

Group	Discrete subgroup	Order	Action Mechanism
$SO(3)$	rotational (chiral)	$ G $	$x \rightarrow \tilde{R}xR$
$O(3)$	reflection (full/Coxeter)	$2 G $	$x \rightarrow \pm \tilde{A}xA$
$Spin(3)$	binary	$2 G $	$(R_1, R_2) \rightarrow R_1 R_2$
$Pin(3)$	pinory (?)	$4 G $	$(A_1, A_2) \rightarrow A_1 A_2$

- e.g. the **chiral icosahedral** group has 60 elements, encoded by 120 spinors, which form the **binary icosahedral** group
- together with the **inversion/pseudoscalar**  $I$  this gives 60 rotations and 60 rotoinversions, i.e. the **full icosahedral** group  $H_3$  in 120 elements doubly covered by 240 pinors



# Representations from Clifford multivector groups

- The usual picture of **orthogonal transformations** on an  $n$ -dimensional vector space is via  $n \times n$  **matrices** acting on vectors, immediately making connections with **representations = matrices satisfying the group multiplication laws**.
- **Easy to construct representations** with (s)pinors in the  $2^n$ -dimensional Clifford algebra as **reshuffling components**.
- Spinors leave the **original**  $n$ -dimensional **vector** space invariant, **reshuffle** the components of the **vector**.
- But can also consider various representation matrices acting on **different subspaces** of the Clifford algebra.

# Representations from Clifford multivector groups – trivial, parity, rotation representations

- The **scalar** subspace is **one-dimensional**.  $\tilde{R}1R = \tilde{R}R = 1$  gives the **trivial representation**, and likewise pinors  $A$  give the **parity**.
- The double-sided action  $\tilde{R}xR$  of spinors  $R$  on a **vector**  $x$  in the  $n$ -dimensional vector space gives an  $n \times n$ -dimensional representation, which is just the usual **rotation matrices**.
- E.g.  $e_1e_2$  acting on  $x = x_1e_1 + x_2e_2 + x_3e_3$  gives  $e_2e_1xe_1e_2 = -x_1e_1 - x_2e_2 + x_3e_3$  which could also be expressed as 
$$\begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -x_1 \\ -x_2 \\ x_3 \end{pmatrix}$$
- If the spinors were acting as  $Rx\tilde{R}$  would give a **potentially different representation**.

# Characters, their norm, and the Frobenius-Schur indicator

- **Similarity** transformed representations are also good representations, but are not fundamentally different: they are **equivalent**.
- So want a measure for a representation that is **invariant** under similarity transformations, e.g. the **trace** aka the **character**  $\chi$  of a matrix
- A **class function** i.e. the same within a conjugacy class because of the cyclicity of the trace
- The **character norm**  $||\chi||^2 := \frac{1}{|G|} \sum_{g \in G} |\chi(g)|^2$
- The **Frobenius-Schur indicator**  $v := \frac{1}{|G|} \sum_{g \in G} \chi(g^2)$

# Real representations of real, complex, and quaternionic type

- $||\chi||^2 = \frac{1}{|G|} \sum_{g \in G} |\chi(g)|^2 = 1$ : representation of **real** type
- $||\chi||^2 = \frac{1}{|G|} \sum_{g \in G} |\chi(g)|^2 = 2$ : representation of **complex** type
- $||\chi||^2 = \frac{1}{|G|} \sum_{g \in G} |\chi(g)|^2 = 4$ : representation of **quaternionic** type
- Theorem: A complex representation is irreducible if and only if  $||\chi||^2 = 1$ .
- Theorem: A **real** representation is **irreducible** if and only if  $||\chi||^2 + \nu(\chi) = 2$ , e.g.  $4 - 2 = 2$  or  $1 + 1 = 2$ .

## Representations from Clifford multivector groups – $8 \times 8$ and $4 \times 4$ (whole algebra / even subalgebra)

- Rather than restricting oneself to the  $n$ -dimensional vector space, one can also define representations by  $2^n \times 2^n$ -matrices acting on the **whole** Clifford algebra, i.e. any element acting on an arbitrary element, e.g. here  $8 \times 8$ .
- Likewise, one can define  $2^{(n-1)} \times 2^{(n-1)}$ -dimensional spinor representations as acting on the **even subalgebra**.
- 3D spinors have **components** in  $(1, e_1 e_2, e_2 e_3, e_3 e_1)$ , **multiplication** with another spinor e.g.  $e_1 e_2$  will **reshuffle** these components  $(e_1 e_2, -1, -e_3 e_1, e_2 e_3)$
- This **reshuffling** can therefore be described by a  $4 \times 4$ -matrix.

## $4 \times 4$ – explicit example: $A_1^3$

- E.g.  $\boxed{\pm e_1, \pm e_2, \pm e_3}$  give the 8 spinors  
 $\boxed{\pm 1, \pm e_1 e_2, \pm e_2 e_3, \pm e_3 e_1}$ , or  $(\pm 1, 0, 0, 0)$  (8 permutations)
- $\|\chi\|^2 = 32/8 = 4$ ,  $v = -2$  and  $\|\chi\|^2 + v = 2$  i.e. **real**  
**irreducible of quaternionic type**

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix},$$

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}$$

# Character table of $Q$

$Q$	1	-1	$\pm e_1 e_2$	$\pm e_2 e_3$	$\pm e_3 e_1$
1	1	1	1	1	1
$1'$	1	1	-1	-1	1
$1''$	1	1	-1	1	-1
$1'''$	1	1	1	-1	-1
2	2	-2	0	0	0
$4_H$	4	-4	0	0	0

## $4 \times 4$ – explicit example: $A_3$

- As a set of **vectors** in 4D, they are  $(\pm 1, 0, 0, 0)$  (8 permutations) ,  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  (16 permutations)
- Conjugacy classes:  
 $1 \cdot 4^2 + 1 \cdot (-4)^2 + 6 \cdot 0^2 + 8 \cdot 2^2 + 8 \cdot (-2)^2 = 32 + 32 + 32 = 96$
- $||\chi||^2 = 96/24 = 4$ ,  $\nu = -2$  and  $||\chi||^2 + \nu = 2$  i.e. **real irreducible of quaternionic type.**



## $3 \times 3$ – explicit example: $H_3$

- Icosahedral spinors are

$(\pm 1, 0, 0, 0)$  (8 permutations),  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  (16 permutations)

$\frac{1}{2}(0, \pm 1, \pm \sigma, \pm \tau)$  (96 even permutations),

- E.g. the rotation matrices corresponding to  $\alpha_1 \alpha_2$  and  $\alpha_2 \alpha_3$  via  $\tilde{R} \times R$  are

$$\frac{1}{2} \begin{pmatrix} \tau & \tau - 1 & -1 \\ 1 - \tau & -1 & -\tau \\ -1 & \tau & 1 - \tau \end{pmatrix} \text{ and } \frac{1}{2} \begin{pmatrix} \tau & 1 - \tau & -1 \\ 1 - \tau & 1 & -\tau \\ 1 & \tau & \tau - 1 \end{pmatrix}.$$

The characters  $\chi(g)$  are obviously 0 and  $\tau$

- $\|\chi\|^2 = 120/120 = 1$ ,  $\nu = 1$  and  $\|\chi\|^2 + \nu = 2$  i.e. real irreducible of real type

## $3 \times 3$ – explicit example: $H_3$ other way

- If the spinors were acting as  $R\tilde{x}\tilde{R}$ , then

$$\frac{1}{2} \begin{pmatrix} \tau & 1-\tau & -1 \\ \tau-1 & -1 & \tau \\ -1 & -\tau & 1-\tau \end{pmatrix} \text{ and } \frac{1}{2} \begin{pmatrix} \tau & 1-\tau & 1 \\ 1-\tau & 1 & \tau \\ -1 & -\tau & \tau-1 \end{pmatrix},$$

with the same characters as before. Swapping the action of the spinor can change the representation.

## $4 \times 4$ – explicit example: $H_3$

- Spinors  $\alpha_1 \alpha_2$  and  $\alpha_2 \alpha_3$  multiplying a generic spinor  $R = a_0 + a_1 e_2 e_3 + a_2 e_3 e_1 + a_3 e_1 e_2$  from the left reshuffles the components  $(a_1, a_2, a_3, a_0)$  with the matrices given as

$$\frac{1}{2} \begin{pmatrix} -1 & \tau - 1 & 0 & -\tau \\ 1 - \tau & -1 & -\tau & 0 \\ 0 & \tau & -1 & \tau - 1 \\ \tau & 0 & 1 - \tau & -1 \end{pmatrix}, \frac{1}{2} \begin{pmatrix} -\tau & 0 & 1 - \tau & -1 \\ 0 & -\tau & -1 & \tau - 1 \\ \tau - 1 & 1 & -\tau & 0 \\ 1 & 1 - \tau & 0 & -\tau \end{pmatrix},$$

with characters  $-2$  and  $-2\tau$ .

## $4 \times 4$ – explicit example $H_3$ : quaternionic type

- 120  $4 \times 4$  matrices – 9 conjugacy classes, with pairs that have  $\pm 2\chi_3$  so gives **4 times** that of the  $3 \times 3$  case
- $|G| \cdot ||\chi||^2 = 1 \cdot 4^2 + 1 \cdot (-4)^2 + 12 \cdot (-2\tau)^2 + 12 \cdot (2\tau)^2 + 12 \cdot (-2\sigma)^2 + 12 \cdot (2\sigma)^2 + 20 \cdot (-2)^2 + 20 \cdot (2)^2 + 30 \cdot 0^2 = \mathbf{480}$
- $||\chi||^2 = 480/120 = \mathbf{4}$ ,  $\nu = \mathbf{-2}$  and  $||\chi||^2 + \nu = \mathbf{2}$  i.e. **real irreducible of quaternionic type**

# Character table of $I = A_5$

$I$	1	$20C_3$	$15C_2$	$12C_5$	$12C_5^2$
1	1	1	1	1	1
3	3	0	-1	$\tau$	$\sigma$
$\bar{3}$	3	0	-1	$\sigma$	$\tau$
4	4	1	0	-1	-1
5	5	-1	1	0	0

# Character table of $2/$

$I$	1	$20C_3$	$30C_2$	$12C_5$	$12C_5^2$	$-1$	$-20C_3$	$-12C_5$	$-12C_5^2$
1	1	1	1	1	1	1	1	1	1
3	3	0	-1	$\tau$	$\sigma$	3	0	$\tau$	$\sigma$
$\bar{3}$	3	0	-1	$\sigma$	$\tau$	3	0	$\sigma$	$\tau$
4	4	1	0	-1	-1	4	1	-1	-1
5	5	-1	1	0	0	5	-1	0	0
2	2	-1	0	$-\sigma$	$-\tau$	-2	1	$\sigma$	$\tau$
2	2	-1	0	$-\tau$	$-\sigma$	-2	1	$\tau$	$\sigma$
4	4	1	0	-1	-1	-4	-1	1	1
6	6	0	0	1	1	-6	0	-1	-1
$4_H$	4	-2	0	$-2\tau$	$-2\sigma$	-4	2	$2\tau$	$2\sigma$
$4_{\tilde{H}}$	4	-2	0	$-2\sigma$	$-2\tau$	-4	2	$2\sigma$	$2\tau$

## A general construction of representations of quaternionic type – canonical representations

- It had so far been **overlooked** that there is a **systematic construction** of representations of **quaternionic type** for 3D polyhedral groups
- This is simply due to the fact that the **spinors** in 3D provide a realisation of the **quaternions**
- Therefore spinors provide 4x4 representations of quaternionic type for **all** (though limited number of) possible groups
- However, they are **canonical** for a choice of 3D **simple roots**, i.e. there is a preferred amongst all similarity transformed versions
- These **simple roots** also determine the 3x3 **rotation** matrices and their **reversed** representations in a similar **canonical** way

## Characters in general

- For a **general spinor**  $R = a_0 + a_1 e_2 e_3 + a_2 e_3 e_1 + a_3 e_1 e_2$  one has **3D character**  $\chi = 3a_0^2 - a_1^2 - a_2^2 - a_3^2$  and **representation**

$$\frac{1}{2} \begin{pmatrix} a_0^2 + a_1^2 - a_2^2 - a_3^2 & -2a_0 a_3 + 2a_1 a_2 & 2a_0 a_2 + 2a_1 a_3 \\ 2a_0 a_3 + 2a_1 a_2 & a_0^2 - a_1^2 + a_2^2 - a_3^2 & -2a_0 a_1 + 2a_2 a_3 \\ -2a_0 a_2 + 2a_1 a_3 & 2a_0 a_1 + 2a_2 a_3 & a_0^2 - a_1^2 - a_2^2 + a_3^2 \end{pmatrix}$$

- and the **4D rep and character** are

$$\begin{pmatrix} a_0 & a_3 & -a_2 & a_1 \\ -a_3 & a_0 & a_1 & a_2 \\ a_2 & -a_1 & a_0 & a_3 \\ -a_1 & -a_2 & -a_3 & a_0 \end{pmatrix}, \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_0 \end{pmatrix} \text{ and } \chi = 4a_0.$$

- Characters** of the representations are **all** determined by the **spinor**!



- 1 Polyhedral groups, Platonic solids and root systems
- 2 Reflection groups with Clifford algebras
  - A Clifford way of doing orthogonal transformations
  - The geometry of the Coxeter plane
  - Root system induction and ADE correspondences
  - Representations from multivector groups
  - Conformal, modular and braid groups
- 3 Conclusions

# Clifford Algebra and orthogonal transformations

- **Inner product** is symmetric part  $a \cdot b = \frac{1}{2}(ab + ba)$
- Reflecting  $a$  in  $b$  is given by  $a' = a - 2(a \cdot b)b = -bab$  ( $b$  and  $-b$  **doubly cover** the same reflection)
- Via **Cartan-Dieudonné** theorem any orthogonal (/conformal/modular) transformation can be written as **successive reflections**

$$x' = \pm n_1 n_2 \dots n_k x n_k \dots n_2 n_1 = \pm A x \tilde{A}$$

- The conformal group  $C(p, q) \sim SO(p+1, q+1)$  so can use these for **translations, inversions** etc as well

# Conformal Geometric Algebra

- Go to  $e_1, e_2, e, \bar{e}$ , with  $e_0^2 = 1, e_i^2 = -1, e^2 = 1, \bar{e}^2 = -1$
- Define two **null** vectors  $n \equiv e + \bar{e}, \bar{n} \equiv e - \bar{e}$
- Can **embed** the 2D vector  $x = x^\mu e_\mu = xe_1 + ye_2$  as a **null vector in 4D** (also normalise  $F(x) \cdot e = -1$ )

$$F(x) = \frac{1}{\lambda^2 - x^2} (x^2 n + 2\lambda x - \lambda^2 \bar{n})$$


- So neat thing is that **conformal transformations** are now done by **rotors** (except inversion which is a reflection) – distances are given by **inner products**

## Conformal Transformations in CGA

$$F(x) = \frac{1}{\lambda^2 - x^2}(x^2 n + 2\lambda x - \lambda^2 \bar{n})$$

- **Reflection:** spacetime  $F(-axa) = -\mathbf{a}F(x)\mathbf{a}$
- **Rotation:** spacetime  $F(Rx\tilde{R}) = RF(x)\tilde{R}$ ,  $R = \exp(\frac{\mathbf{a}\mathbf{b}}{2\lambda})$
- **Translation:**  $F(x+a) = R_T F(x)\tilde{R}_T$  for  $R_T = \exp(\frac{\mathbf{n}\mathbf{a}}{2\lambda}) = 1 + \frac{\mathbf{n}\mathbf{a}}{2\lambda}$
- **Dilation:**  $F(e^\alpha x) = R_D F(x)\tilde{R}_D$  for  $R_D = \exp(\frac{\alpha}{2\lambda} \mathbf{e}\bar{\mathbf{e}})$
- **Inversion:** Reflection in extra dimension  $\mathbf{e}$ :  $F(\frac{x}{x^2}) = -\mathbf{e}F(x)\mathbf{e}$   
sends  $n \leftrightarrow \bar{n}$
- **Special conformal transformation:**  $F(\frac{x}{1+ax}) = R_S F(x)\tilde{R}_S$  for  
 $R_S = R_I R_T R_I$

# Modular group



- Modular generators:  $T : \tau \rightarrow \tau + 1$ ,  $S : \tau \rightarrow -1/\tau$
- $\langle S, T | S^2 = I, (ST)^3 = I \rangle$  CGA rotor version:  $R_Y X \tilde{R}_Y$
- CGA:  $T_X = \exp\left(\frac{ne_1}{2}\right) = 1 + \frac{ne_1}{2}$  and  $S_X = e_1 e$  (slight issue of complex structure  $\tau =$  complex number, not vector in the 2D real plane so map  $e_1 : x_1 e_1 + x_2 e_2 \leftrightarrow x_1 + x_2 e_1 e_2 = x_1 + ix_2$ )
- $(S_X T_X)^3 = -1$  and  $S_X^2 = -1$
- So a 3-fold and a 2-fold rotation in conformal space

## Braid group

- $(S_X T_X)^3 = -1$  and  $S_X^2 = -1$  is inherently **spinorial**
- Of course Clifford construction gives a **double cover**
- The **braid group** is a double cover
- So **Clifford** construction gives the **braid group double cover** of the **modular group**
- $\sigma_1 = \tilde{T}_X = \exp(-n\bar{e}_1/2)$  and  $\sigma_2 = T_X S_X T_X = \exp(-\bar{n}e_1/2)$  satisfying  $\sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2 (= S_X)$
- Nice **symmetry** between the roles of the **point at infinity** and the **origin**
- Might not be known? **Spinorial techniques** might make awkward **modular transformations** more tractable?

# Conclusions

- Clifford algebra provides a very **general** way of doing reflection **group** theory (Cartan-Dieudonné)
- Construction of the **exceptional root systems** from 3D root systems
- More **geometric** approach to the geometry of the **Coxeter plane, degrees and exponents**
- Geometry of 3D space **systematically** and **canonically** gives representations of 4D root systems in terms of **quaternions** and polyhedral **representations of quaternionic type** (among others)

# Conclusions

Thank you!



## Quaternion groups via the geometric product

- The 8 quaternions of the form  $(\pm 1, 0, 0, 0)$  and permutations are the **Lipschitz units**, the **quaternion group** in 8 elements.
- The 8 Lipschitz units together with  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  are the **Hurwitz units**, the **binary tetrahedral group** of order 24.  
Together with the 24 'dual' quaternions of the form  $\frac{1}{\sqrt{2}}(\pm 1, \pm 1, 0, 0)$ , they form the **binary octahedral group** of order 48.
- The 24 Hurwitz units together with the 96 unit quaternions of the form  $(0, \pm \tau, \pm 1, \pm \sigma)$  and even permutations, are called the **Icosians**. The icosian group is isomorphic to the **binary icosahedral group** with 120 elements.
- The unit spinors  $\{1; e_2 e_3; e_3 e_1; e_1 e_2\}$  of  $\text{Cl}(3)$  are isomorphic to the **quaternion algebra**  $\mathbb{H}$ .

## $H_4$ from icosahedral spinors

- The  $H_3$  root system has 30 **roots** e.g. simple roots  $\alpha_1 = e_2, \alpha_2 = -\frac{1}{2}((\tau-1)e_1 + e_2 + \tau e_3)$  and  $\alpha_3 = e_3$ .
- The subgroup of **rotations** is  $A_5$  of order **60**
- These are doubly covered by **120** spinors of the form  $\alpha_1 \alpha_2 = -\frac{1}{2}(1 - (\tau-1)e_1 e_2 + \tau e_2 e_3)$ ,  $\alpha_1 \alpha_3 = e_2 e_3$  and  $\alpha_2 \alpha_3 = -\frac{1}{2}(\tau - (\tau-1)e_3 e_1 + e_2 e_3)$ .
- As a set of **vectors** in 4D, they are

$(\pm 1, 0, 0, 0)$  (8 permutations) ,  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  (16 permutations)

$\frac{1}{2}(0, \pm 1, \pm \sigma, \pm \tau)$  (96 even permutations) ,

which are precisely the 120 roots of the  **$H_4$  root system**.

# Systematic construction of the polyhedral groups


- Multiplying together root vectors in the Clifford algebra gave a **systematic** way of constructing the **binary polyhedral** groups as 3D spinors = **quaternions**.
- The 6/12/18/30 **roots** in  $A_1 \times A_1 \times A_1 / A_3 / B_3 / H_3$  generate 8/24/48/120 **spinors**.
- The **discrete spinor group** is isomorphic to the **quaternion** group  $Q$  / **binary tetrahedral** group  $2T$  / **binary octahedral** group  $2O$  / **binary icosahedral** group  $2I$ ).

$A_1^3$	$A_3$	$B_3$	$H_3$
$A_1^4$	$D_4$	$F_4$	$H_4$

# Quaternionic representations of 3D and 4D Coxeter groups

- Groups  $E_8$ ,  $D_4$ ,  $F_4$  and  $H_4$  have representations in terms of **quaternions**
- **Extensively used** in the high energy physics/quasicrystal/Coxeter/polytope literature and thought of as deeply significant, though not really clear why
- e.g.  $H_4$  consists of 120 elements of the form  $(\pm 1, 0, 0, 0)$ ,  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  and  $(0, \pm \tau, \pm 1, \pm \sigma)$
- Seen as remarkable that the **subset of the 30 pure quaternions** is a realisation of  $H_3$  (**a sub-root system**)
- Similarly,  $B_3$  and  $A_1 \times A_1 \times A_1$  have representations in terms of **pure quaternions**
- Clifford provides a **much simpler geometric explanation**

# Quaternionic representations in the literature



Pure quaternions = Hodge dualised **root vectors**

Quaternions = **spinors**

# Demystifying Quaternionic Representations

- Pure quaternion subset of 4D groups only gives 3D group if the 3D group contains the inversion/pseudoscalar /
- e.g. does not work for the tetrahedral group  $A_3$ , but  $A_3 \rightarrow D_4$  induction still works, with the central node essentially 'spinorial'
- In fact, it goes the other way around: the 3D groups induce the 4D groups via spinors
- The rank-4 groups are also generated (under quaternion multiplication) by two quaternions we can identify as  $R_1 = \alpha_1 \alpha_2$  and  $R_2 = \alpha_2 \alpha_3$
- Can see these are 'spinor generators' and how they don't really contain any more information/roots than the rank-3 groups alone

## Quaternions vs Clifford versors

- **Sandwiching** is often seen as particularly nice feature of the **quaternions giving rotations**
- This is actually a **general feature** of Clifford algebras/versors **in any dimension**; the isomorphism to the **quaternions** is **accidental** to 3D
- However, the **root system** construction does not necessarily generalise
- 2D generalisation merely gives that  $I_2(n)$  is **self-dual**
- **Octonionic** generalisation just induces two copies of the above 4D root systems, e.g.  $A_3 \rightarrow D_4 \oplus D_4$
- Recently constructed  $E_8$  from the **240** pinors doubly covering 120 elements of  $H_3$  in  $2^3 = 8$ -dimensional 3D Clifford algebra